

Double parton distributions in the leading logarithm approximation of perturbative QCD

A.M. Snigirev

M.V.Lomonosov Moscow State University, D.V.Skobeltsyn Institute of Nuclear Physics

119992, Vorobiev Gory, Moscow, Russia

E-mail: snigirev@lav01.sinp.msu.ru

Abstract

Recent CDF measurements of the inclusive cross section for a double parton scattering attach a great importance to any theoretical calculations of two-particle distribution functions. Using a parton interpretation of the leading logarithm diagrams of perturbative QCD theory, generalized Lipatov-Altarelli-Parisi-Dokshitzer equations for the two-parton distributions are re-obtained. The solutions of these equations are not at all the product of two single-parton distributions what is usually applied to the current analysis as ansatz.

PACS: 12.38.Mh, 24.85.+p, 25.75.+r

Keywords: many parton distributions, leading logarithm approximation

The CDF Collaboration has recently measured a large number of double parton scatterings [1]. Thus new and complementary information on the structure of the proton can be obtained by identifying and analyzing events in which two parton-parton hard scatterings take place within one $p\bar{p}$ collision. This process, double parton scattering, provides information on both the spatial distribution of partons within the proton, and possible parton-parton correlations. Both the absolute rate for the double parton process and any dynamics that correlations may introduce are therefore of interest. The theoretical estimations of the effect under consideration have been done in a number of works [2, 3, 4, 5, 6, 7, 8].

For instance, the differential cross section for the four-jet process (due to the simultaneous interaction of two parton pairs) is given by [4, 5]

$$d\sigma = \sum_{q/g} \frac{d\sigma_{12} d\sigma_{34}}{\sigma_{eff}} D_p(x_1, x_3) D_{\bar{p}}(x_2, x_4), \quad (1)$$

$d\sigma_{ij}$ stands for the two-jet cross section. The factor σ_{eff} in the denominator represents the total inelastic cross section which is an estimate for the size of hadron, $\sigma_{eff} \simeq \pi r_p^2$. Based on the simple "hard sphere" model of proton structure, the expected value for σ_{eff} is 11 mb and is consistent with CDF measured value of $(14.5 \pm 1.7^{+1.7}_{-2.3})$ mb [1]. The two-parton distributions are supposed to be the product of two single-parton distributions times a momentum conserving phase space factor

$$D_p(x_i, x_j) = D_p(x_i, Q^2) D_p(x_j, Q^2) (1 - x_i - x_j), \quad (2)$$

where $D_p(x_i, Q^2)$ are the single quark/gluon momentum distributions at the scale Q^2 (determined by a hard process).

The main purpose of this Letter is to analyze the status of the factorization ansatz (2) for many parton distributions in the perturbative QCD theory.

Here one should note that the generalized Lipatov-Altarelli-Parisi-Dokshitzer equations for many parton distribution functions have been derived for the first time in refs. [9, 10] within the leading logarithm approximation of QCD using a method by Lipatov [11]. Under certain initial conditions these equations lead to the solutions, which are identical with the jet calculus rules proposed for multiparton fragmentation functions by Konishi-Ukawa-Veneziano [12]. Because of a very old affair it is necessary to recall some features of that investigation to be clear.

In ref. [13] the structure functions of ep scattering and e^+e^- annihilation were calculated in the leading logarithmic approximation for vector and pseudoscalar theories. Similar calculations in QCD were made in [14]. Lipatov shown [11] that the results of these calculations admit a simple interpretation in the framework of the parton model with a variable cut-off parameter $\Lambda \sim Q^2$ with respect to the transverse momenta, and derived an equation for the scaling

violation of the parton distribution $D_i^j(x, \Lambda)$ inside a dressed quark or gluon, which is in fact equivalent to the one proposed by Altarelli and Parisi [15], within the difference that it was not applied by Lipatov to QCD.

Calculating ladder graph contributions to the structure functions it is convenient to use Sudakov's technique [16]:

$$k = \alpha q' + \beta p + k_\perp, \quad q' = q - \frac{q^2}{s}p, \quad q'^2 = 0, \quad pq' = \frac{1}{2}s, \quad (3)$$

q is the momentum of the virtual photon, p is the momentum ($p^2 = 0$) of the considered quark or gluon. In deep inelastic processes this constituent is one of the partons found at the reference virtualness $-q_0^2 = \mu^2$ in the target proton. In e^+e^- annihilation it is that final quark or gluon which converts into the detected hadron. q' is a light-like vector combined from p and q . $k_\perp$ is the component of k orthogonal to q' and p . In the next step one takes the residues with respect to α in the Feynman integrals. Now each line r carries only β_r and $k_{\perp r}$. And so one may introduce the amplitudes $\Psi_i^n(\beta_r, k_{\perp r})$ for finding in the dressed constituent i n bare partons of given types and parameters $\beta_r, k_{\perp r}, r = 1, \dots, n$. Analogously $\bar{\Psi}_i^n(\beta_r, k_{\perp r})$ are the probability amplitudes for finding a bare parton of i -type in a state of n dressed partons with Sudakov parameters $\beta_r, k_{\perp r}$ in the case of e^+e^- annihilation. These amplitudes can be calculated as tree diagrams in an old-fashioned perturbation theory and obey the normalization condition

$$1 = z_i + \sum_{n=2}^{\infty} \int \prod_{r=1}^n \frac{d\beta_r}{\beta_r} \theta(\beta_r) d^2 k_{\perp r} |\Psi_i^n|^2 \delta^2(\sum k_{\perp r}) \delta(\sum \beta_r - 1), \quad (4)$$

where z_i is the wave function renormalization constant, to be interpreted as the probability that the constituent i remains in the bare state.

An important feature of the leading logarithm diagrams is the ordering of the lines. The absence of interference-type diagrams allows a classical probability interpretation. The integration region in $k_\perp$ essential in the leading logarithm approximation goes up to $\Lambda = |q^2|, |k_\perp| < \Lambda$. The same cut-off is used for the ultraviolet-divergent z_i . The unrenormalized amplitudes $\Psi_i^n, \bar{\Psi}_i^n$ are connected with the renormalized (cut-off independent) ones $\varphi_i^n, \bar{\varphi}_i^n$ via

$$\begin{aligned} \Psi_i^n &= \varphi_i^n \prod_j z_j^{n_j/2}(\Lambda), \\ \bar{\Psi}_i^n &= \bar{\varphi}_i^n z_i^{1/2}(\Lambda), \end{aligned} \quad (5)$$

n_j being the number of partons j in the state n .

In the normalization condition (4) the left-hand side does not depend on Λ , while on the right-hand side the functions Ψ_i^n and the limits of integration over $k_\perp$ depend on Λ . Differentiating (4) with respect to Λ one obtains (see refs. [11, 9, 10] for details)

$$0 = \Lambda \frac{dz_i}{d\Lambda} z_i^{-1} + \omega_i(g_\Lambda^2), \quad (6)$$

where

$$\omega_i(g_\Lambda^2) = \frac{\partial \Pi_i(\Lambda/k^2)}{\partial \ln(\Lambda/|k^2|)} \Big|_{k^2=\Lambda} = \frac{g_\Lambda^2}{8\pi^2} \bar{\omega}_i, \quad (7)$$

Π_i is the one-loop self-energy part of the constituent i , g_Λ is the running coupling constant. In eq. (7) one differentiates only the upper limit of the loop integral.

The parton distribution function can be defined in this framework as

$$D_i^j(x, \Lambda) = z_i \delta_{ij} \delta(x-1) + \sum_{n=2}^{\infty} \sum_{r(j)} \int \prod_{r=1}^n \frac{d\beta_r}{\beta_r} \theta(\beta_r) d^2 k_{\perp r} \cdot |\Psi_i^n|^2 \delta^2(\sum k_{\perp r}) \delta(\sum \beta_r - 1) \delta(\beta_{r(j)} - x). \quad (8)$$

The second sum $\sum_{r(j)}$ runs over all partons of type j in the state n . The result of differentiation reads

$$\Lambda \frac{dD_i^j(x, \Lambda)}{d\Lambda} = \frac{g_\Lambda^2}{8\pi^2} \sum_{j'} \int_x^1 \frac{dx'}{x'} D_i^{j'}(x', \Lambda) P_{j' \rightarrow j} \left(\frac{x}{x'} \right), \quad (9)$$

where

$$\begin{aligned} \frac{g_\Lambda^2}{8\pi^2} \frac{1}{x} P_{j \rightarrow j_1} \left(\frac{x_1}{x} \right) &= \omega_{j \rightarrow j_1}(x \rightarrow x_1) - \delta_{jj_1} \delta(x - x_1) \omega_j(g_\Lambda^2), \\ \omega_{j \rightarrow j_1} &= \sum_{j_1 \leq j'} \omega_{j \rightarrow j_1 j'} + \sum_{j' \leq j_1} \omega_{j \rightarrow j' j_1}, \quad \omega_j = \sum_{j_1 \leq j_2} \omega_{j \rightarrow j_1 j_2}, \end{aligned} \quad (10)$$

$\omega_{j \rightarrow j_1 j_2}(x \rightarrow x_1) dx_1 d\Lambda / \Lambda$, as usual, is the probability that a parton of type j with the fraction x of the longitudinal momentum decays into two partons of types j_1 and j_2 , one of which has the fractions x_1 of the longitudinal momentum and the transverse momentum Λ . These probabilities are defined as in eq.(7) but with the types of partons and the longitudinal momentum fractions in the loop specified. Also, the eq.(6) was used to derive the evolution equation (9).

By introducing the natural variable

$$t = \frac{1}{2\pi b} \ln \left[1 + \frac{g^2(\mu^2)}{4\pi} b \ln \left(\frac{\Lambda}{\mu^2} \right) \right], \quad b = \frac{33 - 2n_f}{12\pi} \text{ in QCD},$$

one obtains

$$\frac{dD_i^j(x, t)}{dt} = \sum_{j'} \int_x^1 \frac{dx'}{x'} D_i^{j'}(x', t) P_{j' \rightarrow j} \left(\frac{x}{x'} \right). \quad (11)$$

This is an equation in the form of Altarelli and Parisi [15]. It is interesting that expression (10) for the kernels P already includes a regularization at $x \rightarrow x'$, which was introduced in ref. [15] afterwards.

Defining the two-parton distribution function

$$\begin{aligned} D_i^{j_1 j_2}(x_1, x_2, \Lambda) &= z_i \delta_{ij_1} \delta_{j_1 j_2} \delta(x_1 - x_2) \delta(x_1 - 1) - \delta_{j_1 j_2} \delta(x_1 - x_2) D_i^{j_1}(x_1, \Lambda) + \\ &+ \sum_{n=2}^{\infty} \sum_{r(j_1)} \sum_{r(j_2)} \int \prod_{r=1}^n \frac{d\beta_r}{\beta_r} \theta(\beta_r) d^2 k_{\perp r} |\Psi_i^n|^2 \delta^2(\sum k_{\perp r}) \delta(1 - \sum \beta_r) \delta(\beta_{r(j_1)} - x_1) \delta(\beta_{r(j_2)} - x_2), \end{aligned} \quad (12)$$

one obtains by differentiating with respect to Λ

$$\begin{aligned} \frac{dD_i^{j_1 j_2}(x_1, x_2, t)}{dt} = & \sum_{j_1'} \int_{x_1}^{1-x_2} \frac{dx_1'}{x_1'} D_i^{j_1' j_2}(x_1', x_2, t) P_{j_1' \rightarrow j_1} \left(\frac{x_1}{x_1'} \right) + \\ & + \sum_{j_2'} \int_{x_2}^{1-x_1} \frac{dx_2'}{x_2'} D_i^{j_1 j_2'}(x_1, x_2', t) P_{j_2' \rightarrow j_2} \left(\frac{x_2}{x_2'} \right) + \sum_{j'} D_i^{j'}(x_1 + x_2, t) \frac{1}{x_1 + x_2} P_{j' \rightarrow j_1 j_2} \left(\frac{x_1}{x_1 + x_2} \right), \end{aligned} \quad (13)$$

where the kernel

$$\frac{g_\Lambda^2}{8\pi^2} \frac{1}{x} P_{j \rightarrow j_1 j_2} \left(\frac{x_1}{x} \right) = \omega_{j \rightarrow j_1 j_2}^{j_1 \leq j_2}(x \rightarrow x_1) + \omega_{j \rightarrow j_1 j_2}^{j_1 \geq j_2}(x \rightarrow x_1) \quad (14)$$

is defined without δ -function regularization. This is the generalized Lipatov-Altarelli-Parisi-Dokshitzer equation for two parton distributions $D_i^{j_1 j_2}(x_1, x_2, t)$, representing the probability that in a dressed constituent i one finds two bare partons of types j_1 and j_2 with the given momentum fractions x_1 and x_2 . The result for the m -parton functions can be found in ref. [9].

It is readily verified by direct substitution that the solution of eq. (13) can be written via the convolution of single distributions [9, 10]

$$\begin{aligned} D_i^{j_1 j_2}(x_1, x_2, t) = & \sum_{j' j_1' j_2'} \int_0^t dt' \int_{x_1}^1 \frac{dz_1}{z_1} \int_{x_2}^{1-x_1} \frac{dz_2}{z_2} D_i^{j'}(z_1 + z_2, t') \frac{1}{z_1 + z_2} P_{j' \rightarrow j_1' j_2'} \left(\frac{z_1}{z_1 + z_2} \right) D_{j_1'}^{j_1} \left(\frac{x_1}{z_1}, t - t' \right) D_{j_2'}^{j_2} \left(\frac{x_2}{z_2}, t - t' \right). \end{aligned} \quad (15)$$

This coincides with the jet calculus rules [12] proposed originally for the fragmentation functions and is the generalization of well-known Gribov-Lipatov relation installed for single functions [13, 14] (the distribution of bare partons inside a dressed constituent is identical to the distribution of dressed constituents in the fragmentation of a bare parton in the leading logarithm approximation). The equations for the multiparton fragmentation functions are obtained by Lipatov's method in a similar way [10] and beyond the given investigation.

The solution (15) shows that the distribution of partons is *correlated* in the leading logarithm approximation:

$$D_i^{j_1 j_2}(x_1, x_2, t) \neq D_i^{j_1}(x_1, t) D_i^{j_2}(x_2, t). \quad (16)$$

Of course, it is interesting to find out the phenomenological issue of the equations under consideration. It can be done within the well-known factorization of soft and hard stages (physics of short and long distances) [17]. As result the equations (11) and (13) describe the evolution of parton distributions in a hadron with t (Q^2), if one replaces the index i by index h only. However, the initial conditions for new equations at $t = 0$ ($Q^2 = \mu^2$) are unknown a priori and must be introduced phenomenologically or must be extracted from experiments or some models dealing with physics of long distances (at the parton level: $D_i^j(x, t = 0) = \delta_{ij} \delta(x - 1)$;

$D_i^{j_1 j_2}(x_1, x_2, t = 0) = 0$). Nevertheless the solution of eq. (13) with the given initial condition may be written as before via the convolution of single distributions [10]

$$D_h^{j_1 j_2}(x_1, x_2, t) = \sum_{j_1' j_2'} \int_{x_1}^1 \frac{dz_1}{z_1} \int_{x_2}^{1-x_1} \frac{dz_2}{z_2} D_h^{j_1' j_2'}(z_1, z_2, 0) D_{j_1'}^{j_1}(\frac{x_1}{z_1}, t) D_{j_2'}^{j_2}(\frac{x_2}{z_2}, t) + \quad (17)$$

$$\sum_{j_1' j_1' j_2'} \int_0^t dt' \int_{x_1}^1 \frac{dz_1}{z_1} \int_{x_2}^{1-x_1} \frac{dz_2}{z_2} D_h^{j_1' j_2'}(z_1 + z_2, t') \frac{1}{z_1 + z_2} P_{j' \rightarrow j_1' j_2'} \left(\frac{z_1}{z_1 + z_2} \right) D_{j_1'}^{j_1}(\frac{x_1}{z_1}, t - t') D_{j_2'}^{j_2}(\frac{x_2}{z_2}, t - t').$$

The reckoning for unsolved confinement problem (physics of long distances) is unknown two-parton correlation function $D_h^{j_1' j_2'}(z_1, z_2, 0)$ at some scale μ^2 . One can suppose that this function is the product of two single-parton distributions times a momentum conserving factor at this scale μ^2 :

$$D_h^{j_1 j_2}(z_1, z_2, 0) = D_h^{j_1}(z_1, 0) D_h^{j_2}(z_2, 0) \theta(1 - z_1 - z_2), \quad (18)$$

then

$$D_h^{j_1 j_2}(x_1, x_2, t) = \left(D_h^{j_1}(x_1, t) D_h^{j_2}(x_2, t) + \right. \quad (19)$$

$$\left. \sum_{j_1' j_2'} \int_{x_1}^1 \frac{dz_1}{z_1} \int_{x_2}^{1-x_1} \frac{dz_2}{z_2} D_h^{j_1'}(z_1, 0) D_h^{j_2'}(z_2, 0) D_{j_1'}^{j_1}(\frac{x_1}{z_1}, t) D_{j_2'}^{j_2}(\frac{x_2}{z_2}, t) [\theta(1 - z_1 - z_2) - 1] \right) \theta(1 - x_1 - x_2) +$$

$$\sum_{j_1' j_1' j_2'} \int_0^t dt' \int_{x_1}^1 \frac{dz_1}{z_1} \int_{x_2}^{1-x_1} \frac{dz_2}{z_2} D_h^{j_1' j_2'}(z_1 + z_2, t') \frac{1}{z_1 + z_2} P_{j' \rightarrow j_1' j_2'} \left(\frac{z_1}{z_1 + z_2} \right) D_{j_1'}^{j_1}(\frac{x_1}{z_1}, t - t') D_{j_2'}^{j_2}(\frac{x_2}{z_2}, t - t'),$$

where

$$D_h^j(x, t) = \sum_{j'} \int_x^1 \frac{dz}{z} D_h^{j'}(z, 0) D_{j'}^j(\frac{x}{z}, t) \quad (20)$$

is the solution of eq. (11) with the given initial condition $D_h^j(x, 0)$ for parton distributions inside a hadron expressed via distributions at the parton level.

This result (19) is, as matter of fact, the answer to the question set. If the two-parton distributions are factorized at some scale μ^2 then the evolution violates this factorization *inevitably* at any different scale ($Q^2 \neq \mu^2$), apart from the violation due to the kinematic correlations induced by the momentum conservation (given by θ -functions)¹.

For a practical employment it is interesting to know the degree of this violation. It can be done numerically using, for instance, the CTEQ-parametrization [18] for single distributions as an input in eq. (19) and considering the kinematics of some specific process. Partly this problem was investigated theoretically in refs. [10, 19] and for the two-particle correlations of fragmentation functions in ref. [20]. That technique is based on the Mellin transformation of distribution functions like

$$M_h^j(n, t) = \int_0^1 dx x^n D_h^j(x, t). \quad (21)$$

¹Here this is the analog of a momentum conserving phase space factor in eq. (2)

After that the integro-differential equations (11) and (13) become systems of ordinary linear-differential equations of first order with constant coefficients and can be solved explicitly [10, 19]. In order to obtain the distributions in x -representation an inverse Mellin transformation must be performed

$$D_h^j(x, t) = \int \frac{dn}{2\pi i} x^{-n} M_h^j(n, t), \quad (22)$$

where the integration runs along the imaginary axis to the right from all n -singularities. This can be done numerically again. However the asymptotic behaviour can be estimated. Namely, with the growth of t (Q^2) the second term in eq. (17) becomes *dominant* for the finite x_1 and x_2 [19]. Thus the two-parton distribution functions "forget" the initial conditions unknown a priori and the correlations perturbatively calculated appear.

The CDF Collaboration found no evidence for the kinematic correlation between the two scatterings in double parton events. This can mean only that the factorization ansatz (2) is the acceptable approximation at the scale $Q_{CDF} \sim 5$ GeV accessible to CDF measurements ($E_T^{jet}(min) \simeq 5$ GeV [1]). There are no arguments to assert that this ansatz (2) is acceptable at the larger scales of hard processes accessible to LHC measurements. The second term in eqs. (17) and (19) can give the meaning contribution to the cross section of many jet production at LHC energy and influence on the background estimations to the Higgs production.

To summarize, the analysis shows that within the leading logarithm approximation of the perturbative QCD theory and the factorization of physics of short and long distances, the two-parton distribution functions being the product of two single distributions at some reference scale become to be dynamically correlated at any different scale of a hard process. These correlations are perturbatively calculable (19). Of course, in order to be more conclusive one needs to do numerical estimations of this effect that is planning to perform in the future.

Acknowledgements

It is pleasure to thank V.A. Ilyin drawing the attention to the problem of a double parton scattering. Discussions with E.E. Boos, I.P. Lokhtin, S.V. Molodtsov, A.S. Proskuryakov, V.I. Savrin and G.M. Zinovjev are gratefully acknowledged.

References

- [1] F. Abe et al. (CDF Collaboration), Phys. Rev. D 56 (1997) 3811.
- [2] C. Goebel, F. Halzen and D.M. Scott, Phys. Rev. D 22 (1980) 2789.
- [3] N. Paver and D. Treleani, Nuovo Cimento A 70 (1982) 215.
- [4] B. Humpert, Phys. Lett. 131 B (1983) 461.
- [5] B. Humpert and R. Odorico, Phys. Lett. 154 B (1985) 211.
- [6] T. Sjostrand and M. van Zijl, Phys. Rev. D 36 (1987) 2019
- [7] G. Calucci and D. Treleani, Nucl. Phys. Proc. Suppl. 71 (1999) 392.
- [8] A. Del Fabbro and D. Treleani, Phys.Rev. D 61 (2000) 077502;
Nucl. Phys. Proc. Suppl. 92 (2001) 130.
- [9] R. Kirschner, Phys. Lett. 84 B (1979) 266.
- [10] V.P. Shelest, A.M. Snigirev and G.M. Zinovjev, Phys. Lett. 113 B (1982) 325;
Teor. Mat. Fiz. 51 (1982) 317.
- [11] L.N. Lipatov, Yad. Fiz. 20 (1974) 181.
- [12] K. Konishi, A. Ukawa and G. Veneziano, Phys. Lett. 78 B (1978) 243;
Nucl. Phys. B 157 (1979) 45.
- [13] V.N. Gribov and L.N. Lipatov, Yad. Fiz. 15 (1972) 781; 1218.
- [14] Yu.L. Dokshitzer, Zh. Eksp. Teor. Fiz. 73 (1977) 1216.
- [15] G. Altarelli and G. Parisi, Nucl. Phys. B 126 (1977) 298.
- [16] V.V. Sudakov, Zh. Eks. Teor. Fiz. 30 (1956) 87.
- [17] J.C. Collins, D.E. Soper and G. Sterman, in *Perturbative QCD*, edited by A. Mueller (World Scientific, Singapore, 1989).
- [18] H.L. Lai et al. (CTEQ Collaboration), Eur. Phys. J. C 12 (2000) 375.
- [19] V.P. Shelest, A.M. Snigirev and G.M. Zinovjev, JINR Communication, E2-82-194, Dubna (1982); Preprint ITP-83-46E, Kiev (1983), unpublished.
- [20] M.J. Puhala, Phys. Rev. D 22 (1980) 1087.